

References

- [i] Zwally et. al., Mass Gains of the Antarctic Ice Sheet Exceed Losses, SCAR ISMASS 7/14/2012
- [ii] Ridley, *Warming could push Greenland beyond tipping points*, U.K. Met Office.gov.uk, 9/27/2013
- [iii] Goldenberg, *Greenland ice sheet faces 'tipping point in 10 years'*, The Guardian 8/10/2010
- [iv] Zwally et. al., *Greenland Ice Sheet Mass Balance*, J. Glaciology 57: 88-102 (2011)
- [v] According to the UK Met office, the ice volume is $2.85E+6 \text{ km}^3$ and the present melt rate is $195 \text{ km}^3/\text{year}$. That would take 14,600 years. Zwally (NASA) estimates the present rate of loss at $171 \text{ km}^3/\text{year}$, or 16700 years. A couple of millennia difference hardly matters when pondering immediate 21st century mitigation.
- [vi] Scheuchl et. al., *Ice Velocity Change in the Ross and Ronne sectors...*, Cryosphere 6: 1019-1030 (2012)
- [vii] Hulbe et. al., *Recent Changes in the flow of the Ross Ice Shelf*, Earth & Planetary Science Letters 376: 54-62 (2013)
- [viii] NASA.gov/jpl/news/Antarctic-ice-sheet-201450512/#U3PDB8blvu1r
- [ix] Rignot et. al., *Acceleration of Pine Island and Thwaites...*, Annals of Glaciology 34: 189-194 (2002).
- [x] Pritchard et. al., *Antarctic Ice Sheet Loss...*, Nature 484: 502-505 (2012)
- [xi] Medley et. al., *Constraining the ice mass balance...with airborne observations of snow accumulation*, Cryosphere Discuss. 8: 953-998 (2014)